

PERIODIC CLASSIFICATION

1. INTRODUCTION

Periodic table may be defined as the table which classifies all the known elements in accordance with their properties in such a way that elements with similar properties are grouped together in the same vertical column and dissimilar elements are separated from one another.

2. HISTORICAL DEVELOPMENT OF THE PERIODIC TABLE

All earlier attempts of the classification of the elements were based upon their atomic weights.

2.1 Dobereiner's Triads

In 1829, Dobereiner classified certain elements in the groups of three called **triads**. The three elements in a triad had similar chemical properties. When the elements in a triad were arranged in the order of increasing atomic weights, the atomic weight of the middle element was found to be approximately equal to the arithmetic mean of the other two elements.

1. Triad	Iron	Cobalt	Nickel	Mean of 1st and 3rd
At. wt.	55.85	58.93	58.71	Atomic weights are nearly the same
2. Triad	Lithium	Sodium	Potassium	
At. wt.	7	23	39	$\frac{7+39}{2} = 23$
3. Triad	Chlorine	Bromine	Iodine	
At. wt.	35.5	80	127	$\frac{35.5+127}{2} = 81.25$
4. Triad	Calcium	Strontium	Barium	
At. wt.	40	87.5	137	$\frac{40+137}{2} = 88.5$

2.2 Newland's Law of Octaves

In 1865, an English chemist, John Alexander Newlands observed that


When the lighter elements were arranged in order of their increasing atomic weights, the properties of every eighth element were similar to those of the first one like the eighth note of a musical scale. This generalization was named as Newlands's law of octaves.

Element	Li	Be	B	C	N	O	F
At. wt.	7	9	11	12	14	16	19
Element	Na	Mg	Al	Si	P	S	Cl
At. wt.	23	24	27	29	31	32	35.5
Element	K	Ca					
At. wt.	39	40					

2.3 Lothar Meyer's Curve

"Physical properties of elements are periodic functions of their atomic masses."

According to Lothar Meyer, elements having similar properties occupy the similar positions in atomic volume viz atomic mass curve



2.4 Mendeleev's Periodic Law

Mendeleev arranged elements in horizontal rows and vertical columns of a table in order of their increasing atomic weights in such a way that the elements with similar properties occupied the same vertical column or group.

PERIODIC CLASSIFICATION

2.5 Modern Periodic Law

In 1913, the English physicist, Henry Moseley observed regularities in the characteristic X-ray spectra of the elements. A plot of $\sqrt{\nu}$ (where ν is frequency of X-rays emitted) against **atomic number (Z)** gave a straight line and not the plot of $\sqrt{\nu}$ vs atomic mass.

Mendeleev's Periodic Law was, therefore, accordingly modified. This is known as the **Modern Periodic Law** and can be stated as :

The physical and chemical properties of the elements are periodic functions of their atomic numbers.

PERIODIC TABLE OF THE ELEMENTS

GROUP

1

1A

1

1.0079

H

HYDROGEN

2

4.0026

He

HELIUM

3

6.941

Li

LITHIUM

4

9.0122

Be

BERYLLIUM

5

10.811

B

BORON

6

12.011

C

CARBON

7

14.007

N

NITROGEN

8

15.999

O

OXYGEN

9

18.998

F

FLUORINE

10

20.180

Ne

NEON

11

22.990

Na

SODIUM

12

24.305

Mg

MAGNESIUM

13

26.982

Al

ALUMINUM

14

28.086

Si

SILICON

15

30.974

P

PHOSPHORUS

16

32.065

S

SULFUR

17

35.453

Cl

CHLORINE

18

39.948

Ar

ARGON

19

39.098

K

POTASSIUM

20

40.078

Ca

CALCIUM

21

44.956

Sc

SCANDIUM

22

47.867

Ti

TITANIUM

23

50.942

V

Vanadium

24

51.996

Cr

CHROMIUM

25

55.938

Mn

MANGANESE

26

55.845

Fe

IRON

27

58.933

Co

COBALT

28

58.693

Ni

NICKEL

29

63.546

Cu

COPPER

30

65.38

Zn

ZINC

31

69.723

Ga

GALLIUM

32

72.64

Ge

GERMANIUM

33

72.922

As

ARSENIC

34

78.96

Se

SELENIUM

35

79.904

Br

BROMINE

36

83.798

Kr

KRYPTON

37

85.468

Rb

RUBIDIUM

38

87.62

Sr

STRONTIUM

39

88.906

Y

YTRIUM

40

90.224

Zr

ZIRCONIUM

41

92.906

Nb

NIOBIUM

42

95.96

Mo

MOLYBDENUM

43

(93)

Tc

TECHNETIUM

44

101.07

Ru

RUTHENIUM

45

102.91

Rh

RHODIUM

46

106.42

Pd

PALLADIUM

47

107.87

Ag

SILVER

48

112.41

Cd

CADMIUM

49

114.82

In

INDIUM

50

118.71

Sn

TIN

51

121.76

Sb

ANTIMONY

52

127.60

Te

TELLURIUM

53

126.90

I

IODINE

54

131.29

Xe

XENON

55

132.91

Cs

CAESIUM

56

137.33

Ba

BARIUM

57-71

La-Lu

Lanthanide

72

178.49

Hf

HAFNIUM

73

180.95

Ta

TANTALUM

74

183.84

W

TUNGSTEN

75

186.21

Re

RHENIUM

76

193.22

Os

OSMIUM

77

194.22

Ir

IRIDIUM

78

195.08

Pt

PLATINUM

79

196.97

Au

GOLD

80

200.59

Hg

MERCURY

81

204.38

Tl

THALLIUM

82

207.2

Pb

LEAD

83

208.98

Bi

BISMUTH

84

(209)

Po

POLONIUM

85

(210)

At

ASTATINE

86

(222)

Rn

RADON

87

(223)

Fr

FRANCIUM

88

(226)

Ra

RADIUM

89-103

Ac-Lr

Actinide

104

(267)

Rf

RUTHERFORDIUM

105

(268)

Db

DBYLIUM

106

(271)

Sg

SEABORGIUM

107

(272)

Bh

BOHRNIUM

108

(277)

Hs

HASSIUM

109

(276)

Mt

MEITNERIUM

110

(281)

Ds

DARMSTADTIUM

111

(283)

Rg

ROSGENIUM

112

(285)

Cn

COOPERNIUM

113

()

Uut

UNUNTRIUM

114

(287)

Uuq

UNUNQUADRIUM

115

()

Uub

UNUNBIUM

116

(291)

Uus

UNUNSEXTIUM

117

()

Uuo

UNUNOCTIUM

118

()

Uuo

UNUNOCTIUM

119

()

Uue

UNUNENNIUM

120

()

Uuh

UNUNHECTIUM

121

()

Uui

UNUNIUNIUM

122

()

Uuq

UNUNQUADRIUM

123

()

Uub

UNUNBIUM

124

()

Uut

UNUNTRIUM

125

()

Uuq

UNUNQUADRIUM

126

()

Uub

UNUNBIUM

127

()

Uut

UNUNTRIUM

128

()

Uuq

UNUNQUADRIUM

129

()

Uub

UNUNBIUM

130

()

Uut

UNUNTRIUM

131

()

Uuq

UNUNQUADRIUM

132

()

Uub

UNUNBIUM

133

()

Uut

UNUNTRIUM

134

()

Uuq

UNUNQUADRIUM

135

()

Uub

UNUNBIUM

136

()

Uut

UNUNTRIUM

137

()

Uuq

UNUNQUADRIUM

138

()

Uub

UNUNBIUM

139

()

Uut

UNUNTRIUM

140

()

Uuq

UNUNQUADRIUM

141

()

Uub

UNUNBIUM

142

()

Uut

UNUNTRIUM

143

()

Uuq

UNUNQUADRIUM

144

()

Uub

UNUNBIUM

145

()

Uut

UNUNTRIUM

146

()

Uuq

UNUNQUADRIUM

147

()

Uub

UNUNBIUM

148

()

Uut

UNUNTRIUM

149

()

Uuq

UNUNQUADRIUM

150

()

Uub

UNUNBIUM

151

()

Uut

UNUNTRIUM

152

()

Uuq

UNUNQUADRIUM

153

()

Uub

UNUNBIUM

154

()

Uut

UNUNTRIUM

155

()

Uuq

UNUNQUADRIUM

156

()

Uub

UNUNBIUM

157

()

Uut

UNUNTRIUM

158

()

Uuq

UNUNQUADRIUM

159

()

Uub

UNUNBIUM

160

()

Uut

UNUNTRIUM

161

()

Uuq

UNUNQUADRIUM

162

()

Uub

UNUNBIUM

163

()

Uut

UNUNTRIUM

164

()

Uuq

UNUNQUADRIUM

165

()

Uub

UNUNBIUM

166

()

Uut

UNUNTRIUM

167

()

Uuq

UNUNQUADRIUM

168

()

Uub

UNUNBIUM

169

()

Uut

UNUNTRIUM

170

()

Uuq

UNUNQUADRIUM

171

()

Uub

UNUNBIUM

172

()

Uut

UNUNTRIUM

173

()

Uuq

UNUNQUADRIUM

174

()

Uub

UNUNBIUM

175

()

Uut

UNUNTRIUM

176

()

Uuq

UNUNQUADRIUM

177

()

Uub

UNUNBIUM

178

()

Uut

UNUNTRIUM

179

()

Uuq

UNUNQUADRIUM

180

()

Uub

UNUNBIUM

181

()

Uut

UNUNTRIUM

182

()

Uuq

UNUNQUADRIUM

183

()

Uub

UNUNBIUM

184

()

Uut

UNUNTRIUM

185

()

Uuq

UNUNQUADRIUM

186

()

Uub

UNUNBIUM

187

()

Uut

UNUNTRIUM

188

()

Uuq

UNUNQUADRIUM

189

()

Uub

UNUNBIUM

190

()

Uut

UNUNTRIUM

191

()

Uuq

UNUNQUADRIUM

192

()

Uub

UNUNBIUM

193

()

Uut

UNUNTRIUM

194

()

Uuq

UNUNQUADRIUM

195

()

Uub

UNUNBIUM

196

()

Uut

UNUNTRIUM

197

()

Uuq

UNUNQUADRIUM

198

()

Uub

UNUNBIUM

199

()

Uut

UNUNTRIUM

200

()

Uuq

UNUNQUADRIUM

201

()

Uub

UNUNBIUM

202

()

Uut

UNUNTRIUM

203

()

Uuq

UNUNQUADRIUM

204

()

Uub

UNUNBIUM

205

()

Uut

UNUNTRIUM

206

()

Uuq

UNUNQUADRIUM

207

()

Uub

UNUNBIUM

208

()

Uut

UNUNTRIUM

209

()

Uuq

UNUNQUADRIUM

210

()

Uub

UNUNBIUM

211

()

Uut

UNUNTRIUM

212

()

Uuq

UNUNQUADRIUM

213

()

Uub

UNUNBIUM

214

()

Uut

UNUNTRIUM

215

()

Uuq

UNUNQUADRIUM

216

()

Uub

UNUNBIUM

217

()

Uut

UNUNTRIUM

218

()

Uuq

UNUNQUADRIUM

219

()

Uub

UNUNBIUM

220

()

Uut

UNUNTRIUM

221

()

Uuq

UNUNQUADRIUM

222

()

Uub

UNUNBIUM

223

()

Uut

UNUNTRIUM

224

()

Uuq

UNUNQUADRIUM

225

()

Uub

UNUNBIUM

226

()

Uut

UNUNTRIUM

227

()

Uuq

UNUNQUADRIUM

228

()

Uub

UNUNBIUM

229

()

Uut

UNUNTRIUM

230

()

Uuq

UNUNQUADRIUM

231

()

Uub

UNUNBIUM

232

()

Uut

UNUNTRIUM

233

()

Uuq

UNUNQUADRIUM

234

()

Uub

UNUNBIUM

235

()

Uut

UNUNTRIUM

236

()

Uuq

UNUNQUADRIUM

237

()

Uub

UNUNBIUM

238

()

Uut

UNUNTRIUM

239

()

Uuq

UNUNQUADRIUM

240

()

Uub

UNUNBIUM

241

()

Uut

UNUNTRIUM

242

()

Uuq

UNUNQUADRIUM

243

()

Uub

UNUNBIUM

244

()

Uut

UNUNTRIUM

245

()

Uuq

UNUNQUADRIUM

246

()

Uub

UNUNBIUM

247

()

Uut

UNUNTRIUM

248

()

Uuq

UNUNQUADRIUM

249

()

Uub

UNUNBIUM

250

()

Uut

UNUNTRIUM

251

()

Uuq

UNUNQUADRIUM

252

()

Uub

UNUNBIUM

253

()

Uut

UNUNTRIUM

254

()

Uuq

UNUNQUADRIUM

255

()

Uub

UNUNBIUM

256

()

Uut

UNUNTRIUM

257

()

Uuq

UNUNQUADRIUM

258

()

Uub

UNUNBIUM

259

()

Uut

UNUNTRIUM

260

()

Uuq

UNUNQUADRIUM

261

()

Uub

UNUNBIUM

262

()

Uut

UNUNTRIUM

263

()

Uuq

UNUNQUADRIUM

264

()

Uub

UNUNBIUM

265

()

Uut

UNUNTRIUM

266

()

Uuq

UNUNQUADRIUM

267

()

Uub

UNUNBIUM

268

()

Uut

UNUNTRIUM

269

()

Uuq

UNUNQUADRIUM

270

()

Uub

UNUNBIUM

271

()

Uut

UNUNTRIUM

272

()

Uuq

UNUNQUADRIUM

273

()

Uub

UNUNBIUM

274

()

Uut

UNUNTRIUM

275

()

Uuq

UNUNQUADRIUM

276

()

Uub

UNUNBIUM

277

()

Uut

UNUNTRIUM

278

()

Uuq

UNUNQUADRIUM

279

()

Uub

UNUNBIUM

280

()

Uut

UNUNTRIUM

281

()

Uuq

UNUNQUADRIUM

282

()

Uub

UNUNBIUM

283

()

Uut

UNUNTRIUM

284

()

Uuq

UNUNQUADRIUM

285

()

Uub

UNUNBIUM

286

()

Uut

UNUNTRIUM

287

()

Uuq

UNUNQUADRIUM

288

()

Uub

UNUNBIUM

289

()

Uut

UNUNTRIUM

290

()

Uuq

UNUNQUADRIUM

291

()

Uub

UNUNBIUM

292

()

Uut

UNUNTRIUM

293

()

Uuq

UNUNQUADRIUM

294

()

Uub

UNUNBIUM

295

()

Uut

UNUNTRIUM

296

()

Uuq

UNUNQUADRIUM

297

()

Uub

UNUNBIUM

298

()

Uut

UNUNTRIUM

299

()

Uuq

UNUNQUADRIUM

300

()

Uub

UNUNBIUM

301

()

Uut

UNUNTRIUM

302

()

Uuq

UNUNQUADRIUM

303

()

Uub

UNUNBIUM

304

()

Uut

UNUNTRIUM

305

()

Uuq

UNUNQUADRIUM

306

()

Uub

UNUNBIUM

307

()

Uut

UNUNTRIUM

308

()

Uuq

UNUNQUADRIUM

309

()

Uub

UNUNBIUM

310

()

Uut

UNUNTRIUM

311

()

Uuq

UNUNQUADRIUM

312

()

Uub

UNUNBIUM

313

()

Uut

UNUNTRIUM

314

()

Uuq

UNUNQUADRIUM

315

()

Uub

UNUNBIUM

316

()

Uut

UNUNTRIUM

317

()

Uuq

UNUNQUADRIUM

318

()

Uub

UNUNBIUM

319

()

Uut

UNUNTRIUM

320

()

Uuq

UNUNQUADRIUM

321

()

Uub

UNUNBIUM

322

()

Uut

UNUNTRIUM

323

()

Uuq

UNUNQUADRIUM

324

()

Uub

UNUNBIUM

325

()

Uut

UNUNTRIUM

326

()

Uuq

UNUNQUADRIUM

327

()

Uub

UNUNBIUM

328

()

Uut

UNUNTRIUM

329

()

Uuq

UNUNQUADRIUM

330

()

Uub

UNUNBIUM

331

()

Uut

UNUNTRIUM

332

()

Uuq

UNUNQUADRIUM

333

()

Uub

UNUNBIUM

334

()

Uut

UNUNTRIUM

335

()

Uuq

UNUNQUADRIUM

336

()

Uub

UNUNBIUM

337

()

Uut

UNUNTRIUM

338

()

Uuq

UNUNQUADRIUM

339

()

Uub

UNUNBIUM

340

()

Uut

UNUNTRIUM

341

()

Uuq

UNUNQUADRIUM

342

()

Uub

UNUNBIUM

343

()

Uut

UNUNTRIUM

344

()

Uuq

UNUNQUADRIUM

345

()

Uub

UNUNBIUM

346

()

Uut

UNUNTRIUM

347

()

Uuq

UNUNQUADRIUM

348

()

Uub

UNUNBIUM

349

()

Uut

UNUNTRIUM

350

()

Uuq

UNUNQUADRIUM

351

()

Uub

UNUNBIUM

352

()

Uut

UNUNTRIUM

353

()

Uuq

UNUNQUADRIUM

354

()

Uub

UNUNBIUM

355

()

Uut

UNUNTRIUM

356

()

Uuq

UNUNQUADRIUM

357

()

Uub

UNUNBIUM

358

()

Uut

UNUNTRIUM

359

()

Uuq

UNUNQUADRIUM

360

()

Uub

UNUNBIUM

361

()

Uut

UNUNTRIUM

362

()

Uuq

UNUNQUADRIUM

363

()

Uub

UNUNBIUM

364

()

Uut

UNUNTRIUM

365

()

Uuq

UNUNQUADRIUM

366

()

Uub

UNUNBIUM

367

()

Uut

UNUNTRIUM

368

()

Uuq

UNUNQUADRIUM

369

()

Uub

UNUNBIUM

370

()

Uut

UNUNTRIUM

371

()

Uuq

UNUNQUADRIUM

372

()

Uub

UNUNBIUM

373

()

Uut

UNUNTRIUM

374

()

Uuq

UNUNQUADRIUM

375

()

Uub

UNUNBIUM

376

()

Uut

UNUNTRIUM

377

()

Uuq

UNUNQUADRIUM

378

()

Uub

UNUNBIUM

379


()

Uut

UNUNTRIUM

</

Classification of Modern Periodic Table



Nomenclature of elements with Atomic Numbers > 100

The naming of the new elements had been traditionally the privilege of the discoverer and the suggested name was ratified by the IUPAC.

Table : Notation for IUPAC Nomenclature of Elements

Digit	Name	Abbreviation
0	nil	n
1	un	u
2	bi	b
3	tri	t
4	quad	q
5	pent	p
6	hex	h
7	sept	s
8	oct	o
9	enn	e

Table : Nomenclature of Elements with Atomic Number Above 100

Atomic Number	Name	Symbol
101	Unnilunium	Unu
102	Unbibium	Unb
103	Unniltrium	Unt
104	Unnilquadium	Unq
105	Unilpentium	Unp
106	Unnilhexium	Unh
107	Unnilseptium	Uns
108	Unniloctium	Uno
109	Unnilennium	Une
110	Ununillium	Uun

3. PREDICTION OF BLOCK, PERIOD & GROUP

1. What electronic configuration
2. Block - last e^- enters into which orbital
3. Period - Max value of principal quantum number

Group - s block - no. of valence electron

p block - $10 + \text{no. of valence electron}$

d block - $ns + \text{no. of } (n-1) d e^-$

f block - III B


4. PROPERTIES OF AN ELEMENT

4.1 Atomic Radius

We cannot measure the exact size of an isolated atom because its outermost electron have a remote chance of being found quite far from the nucleus. So different types of atomic radius can be used based on the environment of atoms i.e; **covalent radius**, **van der Waals' radius**, **metallic radius**.

4.1.1 Covalent Radius

The half of the distance between the nuclei of two identical atoms joined by single covalent bond in a molecule is known as **covalent radius**.



So covalent radius for A-A

$$r_A = \frac{d_{A-A}}{2}$$


If covalent bond is formed between two different elements then

$$d_{A-B} = r_A + r_B - 0.09 (\chi_A - \chi_B)$$

where χ_A and χ_B are electronegative of A and B

4.1.2 Vander Waal's Radius

It is half of the internuclear distance between adjacent atoms of the two neighbouring molecules in the solid state.




$$r_{\text{vander waal}} = \frac{d_{A-A}}{2}$$

PERIODIC CLASSIFICATION

4.1.3 Metallic Radius (Crystal radius)

It is one-half of the distance between the nuclei of two adjacent metal atoms in the metallic crystal lattice.



So metallic radius for A-A

$$d = r_A + r_A$$

$$r_A = \frac{d}{2}$$

$$* r_{\text{covalent}} < r_{\text{metallic}} < r_{\text{vander waals}} *$$

4.2 Variation of Atomic Radii in the Periodic Table

(a) Variation along a period

In general, the covalent and van der Waals radii decrease with increase in atomic number as we move from left to right in a period.

4.3 Atomic Radii

(a) Variation along a period

It is because within the period the outer electrons are in the same valence shell & the effective nuclear charge increases as the atomic number increases resulting in the increased attraction of electrons to the nucleus.

(b) Variation along a group

Atomic radius in a group increases as the atomic number increases. It is because within the group, the principal quantum number (n) increases and the valence electrons are farther from the nucleus.

(c) Ionic Radius

The removal of an electron from an atom results in the formation of a **cation**, whereas gain of an electron leads to an **anion**.

In general, the ionic radii of elements exhibit the same trend as the atomic radii. A cation is **smaller** than its parent atom because it has fewer electrons while its nuclear charge remains the same. The size of an anion will be **larger** than that of the parent atom because the addition of one or more electrons would result in increased repulsion among the electrons and a decrease in effective nuclear charge. For example, the ionic radius of fluoride ion (F^-) is 136 pm whereas the atomic radius of fluorine is only 64 pm. On the other hand, the atomic radius of sodium is 186 pm compared to the ionic radius of 95 pm for Na^+ .


(d) Isoelectronic Species

Isoelectronic species are those which have the same number of electrons. For example, O^{2-} , F^- , Na^+ and Mg^{2+} have the same number of electrons (10). Their radii would be different because of their different nuclear charges. The cation with the greater positive charge will have a smaller radius because of the greater attraction of the electrons to the nucleus. Anion with the greater negative charge will have the larger radius. In this case, the net repulsion of the electrons will outweigh the nuclear charge and the ion will expand in size.

Order of atomic radii is


$$Mg^{2+} < Na^+ < F^- < O^{2-}$$

General Trend :



4.4 Ionization Energy

The minimum amount of energy required to remove the electron from the outermost orbit of an isolated atom in the gaseous state is known as ionization energy.



IE_1 , IE_2 , IE_3 and IE_4 are successive ionization energies.

$$\text{IE}_4 > \text{IE}_3 > \text{IE}_2 > \text{IE}_1$$

$$\text{or } \Delta_1\text{H}_4 > \Delta_1\text{H}_3 > \Delta_1\text{H}_2 > \Delta_1\text{H}_1$$

Variation of Ionisation Energy in Periodic Table

(a) Variation along a period

In a period, the value of ionisation enthalpy increases from left to right with breaks where the atoms have somewhat stable configurations. The observed trends can be easily explained on the basis of increased nuclear charge and decrease in atomic radii. Both the factors increase the force of attraction towards nucleus and consequently, more and more energy is required to remove the electrons and hence, ionisation enthalpies increase.

(b) Variation along a group

On moving the group, the atomic size increases gradually due to an addition of one new principal energy shell at each succeeding element. On account of this, the force of attraction towards the valence electrons decreases and hence the ionisation enthalpy value decreases.

4.5 Units of IE/I.P.

It is measured in units of electron volts (eV) per atom or kilo calories per mole (kcal mol^{-1}) or kilo Joules per mole (kJ mol^{-1}). One electron volt is the energy acquired by an electron while moving under a potential difference of one volt.


$$\begin{aligned} &1 \text{ electron volt (eV) per atom} \\ &= 3.83 \times 10^{-30} \text{ cal per atom} \\ &= 1.602 \times 10^{-19} \text{ J per atom (1 cal} = 4.184 \text{ J)} \\ &= 3.83 \times 10^{-30} \times 6.023 \times 10^{23} \text{ cal mol}^{-1} \\ &= 23.06 \text{ kcal mol}^{-1} \\ &= 1.602 \times 10^{-19} \times 6.023 \times 10^{23} \text{ J mol}^{-1} \\ &= 96.49 \text{ kJ mol}^{-1} \end{aligned}$$

$$\therefore 1 \text{ electron volt (eV) per atom} \\ = 23.06 \text{ kcal mol}^{-1} = 96.49 \text{ kJ mol}^{-1}$$

Important Points


- * Ionization energy increases with decreasing the size of an atom or an ion
- * Ionization energy increases with decreasing screening effect.
- * Ionization energy increases with increasing nuclear charge

- * Ionization energy increases if atom having half filled and fully filled orbitals
- * The penetrating power of orbitals is in the order $s > p > d > f$



Electron Gain Enthalpy

When an electron is added to a neutral gaseous atom (X) to convert it into a negative ion, the enthalpy change accompanying the process is defined as the **Electron Gain Enthalpy** ($\Delta_{\text{eg}}\text{H}$). Electron gain enthalpy provides a measure of the ease with which an atom adds an electron to form anion as represented by



Depending on the element, the process of adding an electron to the atom can be either **endothermic** or **exothermic**. For many elements energy is released when an electron is added to the atom and the electron gain enthalpy is negative. For example, group 17 elements (the **halogens**) have very **high negative electron gain enthalpies** because they can attain stable noble gas electronic configurations by picking up an electron. On the other hand, **noble gases** have large **positive** electron gain enthalpies because the electron has to enter the next higher principal quantum level leading to a very unstable electronic configuration.

Variation of Electron Gain Enthalpy

(a) Variation along a period


Electron gain enthalpy becomes more and more negative from left to right in a period. This is due to decrease in size and increase in nuclear charge as the atomic number increases in a period. Both these factors favour the addition of an extra electron due to higher force of attraction by the nucleus for the incoming electron.

(b) Variation along a group

The electron gain enthalpies, in general, become less negative in going down from top to bottom in a group. This is due to increase in size on moving down a group.

PERIODIC CLASSIFICATION

This factor is predominant in comparison to other factor, i.e., increase in nuclear charge.



4.6 Electronegativity

The tendency of an atom to attract the shared pair of electrons towards itself is known as its **electronegativity**.

According to **Pauling**, the electronegativity of F is 4.0 and electronegativity of other elements can be calculated as

$$(\chi_A - \chi_B) = 0.208 [E_{A-B} - (E_{A-A} \times E_{B-B})^{1/2}]^{1/2}$$


According to **Mulliken**

$$\text{Electronegativity} = \frac{\text{IP} + \text{EA}}{2}$$


(where IP = Ionization potential, EA = Electron affinity)

If IP and EA are taken in electron volt

- * Percentage ionic character = $16(\chi_A - \chi_B) + 3.5(\chi_A - \chi_B)^2$ where χ_A and χ_B are electronegativities of A and B.
- * If the difference in the electronegativities of combining atoms is 1.7, the bond is 50% covalent and 50% ionic.
- * If the difference in electronegativities of oxygen and element is very high the oxide shows a basic character.



The periodic trends of elements in the periodic table



4.7 Periodic Trends in Chemical Properties

4.7.1 Periodicity of Valence or Oxidation States

The electrons present in the outermost shell of an atom are called **valence electrons** and the number of these electrons determine the **valence** or the **valency** of the atom. It is because of this reason that the outermost shell is also called the **valence shell** of the atom and the orbitals present in the valence shell are called **valence orbitals**.

In case of representative elements, the valence of an atom is generally equal to either the number of valence electrons (s- and p-block elements) or equal to eight minus the number of valence electrons.

Group	1	2	13	14	15	16	17	18
Number of valence electrons	1	2	3	4	5	6	7	8
Valence	1	2	3	4	3, 5	2, 6	1, 7	0, 8

In contrast, transition and inner transition elements, exhibit variable valence due to involvement of not only the valence electrons but d- or f-electrons as well. However, their most common valence are 2 and 3.

Let us now discuss periodicity of valence along a period and within a group.

(a) Variation along a period

As we move across a period from left to right, the number of valence electrons increases from 1 to 8. But the valence of elements, w.r.t. H or O first increases from 1 to 4 and then decreases to zero.

In the formation of Na_2O molecule, oxygen being more electronegative accepts two electrons, one from each of the two sodium atoms and thus shows an oxidation state of -2 . On the other hand, sodium with valence shell electronic configuration as $3s^1$ loses one electron to oxygen and is given an oxidation state of $+1$. Thus, the *oxidation state of an element in a given compound may be defined as the charge acquired by its atom on the basis of electronegativity of the other atoms in the molecule.*


(b) Variation within a group

When we move down the group, the number of valence electrons remains the same, therefore, all the elements in a group exhibit the same valence. For example, all the elements of group 1 (alkali metals) have valence one while all the elements of group 2 (alkaline earth metals) exhibit a valence of two.

Noble gases present in group 18 are zerovalent, i.e., their valence is zero since these elements are **chemically inert**.

4.7.2 Anomalous Properties of Second Period Elements

It has been observed that *some elements of the second period show similarities with the elements of the third period placed diagonally to each other, though belonging to different groups*. For example, lithium (of group 1) resembles magnesium (of group 2) and beryllium (of group 2) resembles aluminium (of group 13) and so as. This similarity in properties of elements placed diagonally to each other is called **diagonal relationship**.



The anomalous behaviour is due to their small size, large charge/radius ratio and high electronegativity of the elements. In addition, the first member of group has only four valence orbitals (2s and 2p) available for bonding, whereas the second member of the groups have nine valence orbitals (3s, 3p, 3d). As a consequence of this, the maximum covalency of the first member of each group is 4 (e.g., boron can only form $[\text{BF}_4]^-$, whereas the other members of the groups can expand their valence shell to accommodate more than four pairs of electrons e.g., aluminium forms $[\text{AlF}_6]^{3-}$. Furthermore, the first member of p-block elements displays greater ability to form p π -p π multiple bonds to itself (e.g., $\text{C}=\text{C}$, $\text{C}\equiv\text{C}$, $\text{N}=\text{N}$, $\text{N}\equiv\text{N}$) and to other second period elements (e.g., $\text{C}=\text{O}$, $\text{C}=\text{N}$, $\text{C}\equiv\text{N}$, $\text{N}=\text{O}$) compared to subsequent members of the same groups.


4.7.3 Periodic Trends and Chemical Reactivity

Reactivity of Metals

The reactivity of metals is measured in terms of their tendency to lose electrons from their outermost shell.


In a period

The tendency of an element to lose electrons decreases in going from left to right in a period. So, the reactivity of metals decreases in a period from left to right. For example, the reactivity of third period elements follows the order.



In a group

The tendency to lose electrons increases as we go down a group. So, the reactivity of metals increases down the group. Thus, in group 1, the reactivity follows the order.



— Reactivity increases —>


Reactivity of Non-Metals

The reactivity of a non-metal is measured in terms of its tendency to gain electrons to form an anion.

PERIODIC CLASSIFICATION

In a period

The reactivity of non-metals increases from left to right in a period. During reaction, non-metals tend to form anions. For example, in the second period, the reactivity of non-metals increases in the order.




— Reactivity increases —→

In a group


The reactivity of non-metals in a group decreases as we go down the group. This is because the tendency to accept electrons

decreases down the group. The reactivity of halogens follows the order



— Reactivity decreases —→

The normal oxide formed by the element on extreme left is the **most basic** (e.g., Na_2O) whereas that formed by the element on extreme right is the **most acidic** (e.g., Cl_2O_7). Oxides of elements in the centre are **amphoteric** (e.g., Al_2O_3 , As_2O_3) or **neutral** (e.g., CO , NO , N_2O). Amphoteric oxides behave as acidic with bases and as basic with acids, whereas neutral oxides have no acidic or basic properties.



4.7.4 Inert Pair Effect

In groups 13-16, as we move down the group, the tendency of s-electrons of the valence shell to participate in bond formation decreases. This means that lower oxidation state becomes more stable.

Reason: As we go down these groups, the increased nuclear charge outweighs the effect of the corresponding increase in atomic size. The s-electrons thus become more tightly held (more penetrating) and hence more reluctant to participate in bond formation. Hence, the lower oxidation state becomes more stable.

5. SUMMARY AND IMPORTANT POINTS TO REMEMBER

1. Mendeleev's periodic table was based on atomic masses of the elements. When Mendeleev presented the periodic table, only 63 elements were known. He left 29 places in the table for unknown elements.
2. Modern Mendeleev periodic table is based on atomic numbers of the elements. The modern periodic law is : "**The physical and chemical properties of the elements are periodic function of their atomic numbers**".

The horizontal row in the periodic table is called a **period** and vertical column is called **group**. There are seven periods and nine groups in the modern Mendeleev periodic table.

3. The long or extended form of periodic table consists of seven periods and eighteen vertical columns (groups or families). The elements in a period have same number of energy shells, i.e., principal quantum number (n). These are numbered 1 to 7.

1st period	1s	2 elements
2nd period	2s 2p	8 elements
3rd period	3s 3p	8 elements
4th period	4s 3d 4p	18 elements
5th period	5s 4d 5p	18 elements
6th period	6s 4f 5d 6p	32 elements
7th period	7s 7f 6d 7p	32 elements
Total		*118 elements

At present 114 elements are known.

In a vertical column (group), the elements have similar valence shell electronic configuration and therefore exhibit similar chemical properties.

4. There are four blocks of elements: s-, p-, d- and f-block depending on the orbital which gets the last electron. The general electronic configuration of these blocks are :
 s-block : [Noble gas] $ns^{1 \text{ or } 2}$. However, hydrogen has 1s1 configuration.
 p-block : [Noble gas] $ns^2 np^{1-6}$
 d-block : [Noble gas] $(n-1)d^{1-10} ns^{1 \text{ or } 2}$
 f-block : [Noble gas] $(n-2)f^{1-14} (n-1)d^0 \text{ or } 1 ns^2$

s-block elements occupy IA(1) and IIA(2) groups, i.e., extreme left portion of the periodic table.

p-block elements occupy IIIA(13), IVA(14), VA(15), VIA(16), VIIA(17) and VIIIA(18) groups, i.e., right portion of the periodic table.

d-block elements occupy IIIB(3), IVB(4), VB(5), VIB(6), VIIB(7), VIIIB(8, 9 and 10), IB(11) and IIB(12) groups, i.e., central portion of the periodic table. There are four d-block series, i.e., 3d series, 4d series, 5d series and 6d series, each consisting of ten elements, i.e., in all forty d-block elements are present in periodic table.

f-block elements are accommodated in two horizontal rows below the main periodic table, each row consists of 14 elements, i.e., 28 f-block elements are present in periodic table. The elements in first row are termed 4f-elements or rare earth or lanthanides while the elements of second row are termed 5f-elements or actinides.

5. The elements are broadly divided into three types :
 (i) **Metals** comprise more than 78% of the known elements. s-block, d-block and f-block elements are metals. The higher members of p-block are also metals.
 (ii) **Non-metals** are less than twenty. (C, N, P, O, S, Se, H, F, Cl, Br, I, He, Ne, Ar, Kr, Xe and Rn are non-metals).
 (iii) Elements which lie in the border line between metals and non-metals are called **semimetals** or **metalloids**. B, Si, Ge, As, Sb, Te, Po and At are regarded metalloids.
6. IUPAC given a new scheme for assigning a temporary name to the newly discovered elements. The name is derived directly from the atomic number of the elements. However, IUPAC has accepted the following names of the elements from atomic numbers 104 to 110.

Rutherfordium (Rf), 104	Dubnium (Db), 105	Seaborgium (Sg) 106
Bohrium (Bh), 107	Hassium (Hs), 108	Meitnerium (Mt), 109
Darmstadtium (Ds) 110		

The temporary names of the elements discovered recently are :

Ununium (Uuu), 111	Unubium (Uub) 112
Ununquadium (Uuq) 114	Ununhexium (Uuh) 116

7. The recurrence of similar properties of the elements after certain definite intervals when the elements are arranged in order of increasing atomic numbers in the periodic table is termed **periodicity**. The cause of periodicity is the repetition of similar electronic configuration of the atom in the valence shell after certain definite intervals. These definite intervals are 2, 8, 8, 18, 18 and 32. These are known as magic number. **Periodicity is observed in a number of properties which are directly or indirectly linked with electronic configuration.**
 (i) Effective nuclear charge increases across each period.

PERIODIC CLASSIFICATION

- (ii) Atomic radii generally decrease across the periods.
- (iii) Atomic radii generally increase on moving from top to bottom in the groups.
- (iv) Atomic radius is of three types :
 - (a) **Covalent radius** : It is half of the distance between the centres of the nuclei of two similar atoms joined by a single covalent bond. This is generally used for non-metals.
 - (b) **Crystal or metallic radius** : It is half of the internuclear distance between two nearest atoms in the metallic lattice. It is generally used for metals.
 - (c) **van der Waals' radius** : It is half of the internuclear distance between the nearest atoms belonging to two adjacent molecules in solid state.

van der Waals' radius > Metallic radius > Covalent radius
(for an atom)

- (v) Cations are generally smaller than anions.
- (vi) Cations are smaller and anions are larger than neutral atoms of the elements.

Cation size < Neutral atom size < Anion size

- (vii) Elements of 2nd and 3rd transition series belonging to same vertical columns are similar in size and properties due to **lanthanide contraction**.
- (viii) The first element in each group of the representative elements shows abnormal properties, i.e., differs from other elements of the group because of much smaller size of the atom.
- (ix) The ions having same number of electrons but different nuclear charge are called **isoelectronic ions**.

Examples,

- (a) N^{3-} , O^{2-} , F^- , Na^+ , Mg^{2+} , Al^{3+}
- (b) P^{3-} , S^{2-} , Cl^- , K^+ , Ca^{2+} , Sc^{3+}

In isoelectronic ions, the size decreases if Z/e increases i.e., greater the nuclear charge, smaller is the size of the ion.

- (x) The energy required to remove the most loosely held electron from the gaseous isolated atom is termed **ionisation enthalpy**.
- (xi) Ionisation enthalpy values generally increase across the periods.
- (xii) Ionisation enthalpy values generally decrease down the group.
- (xiii) Removal of electron from filled and half filled shells requires of higher energy. For example, the ionisation enthalpy of nitrogen is higher than oxygen. Be, Mg and noble gases have high values.
- (xiv) Metals have low ionisation enthalpy values while non-metals have high ionisation enthalpy values.
- (xv) Successive ionisation enthalpies of an atom have higher values.
 $IE_1 < IE_2 < IE_3 \dots$
- (xvi) The enthalpy change taking place when an electron is added to an isolated gaseous atom of the element is called **electron**

gain enthalpy. The first electron gain enthalpy of most of the elements is negative as energy is released in the process but the values are positive or near zero in case of the atoms having stable configuration such as Be, Mg, N, noble gases, etc.

- (xvii) Electron gain enthalpy becomes more negative from left to right in a period and less negative from top to bottom in a group.
- (xviii) Successive electron gain enthalpies are always positive.
- (xix) The elements with higher ionisation enthalpy have higher negative electron gain enthalpy.
- (xx) Electronegativity is the tendency of an atom to attract the shared pair of electrons towards itself in a bond.
- (xxi) Electronegativity increases across the periods and decreases down the groups.
- (xxii) Metals have low electronegativities and non-metals have high electronegativities.
- (xxiii) Metallic character decreases across the periods and increases down the group.
- (xxiv) Valence of an element belonging to s- and p- block (except noble gases) is either equal to the number of valence electrons or eight minus number of valence electrons.
- (xxv) The **reducing nature** of the elements decreases across the period while **oxidising nature** increases.
- (xxvi) The **basic character** of the oxides decreases while the **acidic character** increases in moving from left to right in a period.

6. SOME IMPORTANT FACTS ABOUT ELEMENTS

- (i) Bromine is a non-metal which is liquid at room temperature.
- (ii) Mercury is the only metal that is liquid at room temperature.
- (iii) Gallium (m.pt. 29.8°C), caesium (m.pt. 28.5°C) and francium (m.pt. 27°C) are metals having low melting points.
- (iv) Tungsten (W) has the highest melting point (3380°C) among metals.
- (v) Carbon has the highest melting point (4100°C) among non-metals.
- (vi) Oxygen is the most abundant element on the earth.
- (vii) Aluminium is the most abundant metal.
- (viii) Iron is the most abundant transition metal.
- (ix) Highest density is shown by osmium (22.57 g cm^{-3}) or iridium (22.61 g cm^{-3}).
- (x) Lithium is the lightest metal. Its density is 0.54 g cm^{-3} .
- (xi) Silver is the best conductor of electricity.
- (xii) Diamond (carbon) is the hardest natural substance.
- (xiii) Francium has the highest atomic volume.
- (xiv) Boron has the lowest atomic volume.
- (xv) The most abundant gas in atmosphere is nitrogen.
- (xvi) Fluorine is the most electronegative element.
- (xvii) Chlorine has the maximum negative electron gain enthalpy.

- (xviii) Helium has the maximum ionisation enthalpy.
- (xix) Cesium or francium has the lowest ionisation enthalpy.
- (xx) Helium and francium are smallest and largest atoms respectively.
- (xxi) H^- and I^- ions are the smallest and largest anions respectively.
- (xxii) H^+ and Cs^+ ions are the smallest and largest cations respectively.
- (xxiii) Cesium is the most electropositive element.
- (xxiv) Element kept in water is phosphorus, P_4 (white or yellow).
- (xxv) Element kept in kerosene are Na, K, Rb, Cs, etc.
- (xxvi) Iodine is the element which sublimates.
- (xxvii) Hydrogen is the most abundant element in the universe.
- (xxviii) Only ozone is the coloured gas with garlic smell.
- (xxix) Metalloids have electronegativity values closer to 2.0.
- (xxx) First synthetic (i.e., man-made) element is technetium (At. No. 43).
- (xxxi) Most poisonous metal-Plutonium.
- (xxxii) Rarest element in earth's crust-Astatine.
- (xxxiii) The elements coming after uranium are called transuranic elements. The elements with $Z = 104 - 112$, 114 and 116 are called trans-actinides or super heavy elements. All these

elements are synthetic, i.e., man-made elements. These are radioactive elements and not found in nature.

(xxxiv) The elements ruthenium (Ru), germanium (Ge), polonium (Po) and americium (Am) were named in honour of the countries named **Ruthenia (Russia), Germany, Poland and America**, respectively.

(xxxv) The members of the actinide series are radioactive and majority of them are not found in nature.

(xxxvi) The element rutherfordium (Rf, 104) is also called Kurchatovium (Ku) and element dubnium (Db, 105), is also called hahnium.

(xxxvii) Promethium (Pm, 61) a member of lanthanide series is not found in nature. It is a synthetic element.

(xxxviii) Special names are given to the members of these groups in periodic table.

Group 1	or	IA	Alkali metals
Group 2	or	IIA	Alkaline earth metals
Group 15	or	VA	Pnicogens
Group 16	or	VIA	Chalcogens
Group 17	or	VIIA	Halogens
Group 18	or	VIIIA (zero)	Inert or noble gases