


Structure of the Atom

- Atoms are not indivisible and are composed of three fundamental particles. These particles are electrons, protons, and neutrons.
- **Charged particles in Matter**
 - Electrons are negatively-charged particles. They were discovered by J. J. Thomson, by cathode ray experiment.
 - Canal rays are positively charged radiation consisting of protons. Protons are positively-charged particles and were discovered by E. Goldstein.
 - The third fundamental particles present in an atom are neutrons. They are electrically-neutral and were discovered by J. Chadwick.
 - Various models were given to explain the structure of atom.
- **Thomson's atomic model:**
 - Thomson thought that an atom is a sphere of positive charge in which electrons are embedded.
 - An atom as a whole is electrically neutral because the negative and positive charges are equal in magnitude.



- **Rutherford's atomic model:**
 - On the basis of his experiments with alpha rays and gold foil, Rutherford concluded that Thomson's atomic model was incorrect.
 - He proposed an atomic model based on the results of his experiments.
 - In this model, all the positive charges (i.e., protons) were present at the centre of the atom, inside the nucleus, and the electrons were present in circular orbits around the nucleus.
 - He said that the electrons are not at rest and keep moving continuously in these circular orbits.





- He also said that the size of the nucleus is very small as compared to that of the atom.

- **Drawbacks of Rutherford's Model**

- It cannot explain the stability of an atom on the basis of classical mechanics and electromagnetic theory.
- If the electrons were stationary, then the strong electrostatic force of attraction between the dense nucleus and the electrons would pull the electrons towards the nucleus. Thus, it cannot explain the stability of an atom.
- Rutherford's model does not give any idea about the distribution of electrons around the nucleus (i.e., the electronic structure of the atom), and about their energy.
- It cannot explain the atomic spectra.

- **Bohr's atomic model:**

- Neils Bohr proposed that the electrons present around the nucleus revolve in specific orbits called energy levels.
- He also stated that the electrons do not release energy while revolving. Thus, the resulting atom is a stable one.
- The shells in which the electrons are present are known as K, L, M, N, and so on (or 1, 2, 3, 4, and so on), as proposed by Bohr and Bury.
- Each shell contains a specific number of electrons, which can be calculated using the formula $2n^2$.

Atomic Models	
	
Dalton's atomic model	Thomson's atomic model
	
Rutherford's atomic model	Bohr's atomic model

- Valency is defined as the combining capacity of the atom of an element. Valency of an element depends upon the number of electrons present in the outermost shell of its atom.
- **Atomic number** of an element is equal to the number of protons present in the atom and atomic mass is equal to the sum of the number of protons and neutrons present in it.
- **Isotopes** are atoms having the same atomic number and different atomic masses.
- **Isobars** are atoms having the same atomic mass and different atomic numbers.

