

47. The Special Theory of Relativity

Short Answer

1. Question

Answer

The second postulates describe that the speed of light (for medium vacuum) has same value in all inertial frame. Speed of light in any other medium will be less than that of vacuum using Snell's law. Snell's law is given by,

$$v = \frac{c}{\mu}$$

Where,

v=Speed of light in glass

c=Speed of light in vacuum

μ =Refractive index

Thus, it doesn't violate second postulate of the refractive index because the speed of light in glass is not more than that of vacuum.

2. Question

Answer

As the platform is stationary it is considered as the proper frame for this event.

The train travels with a speed which is very less as compared with the speed of light. So, it does not violate the second postulate of relativity.

Thus, both train and platform are proper frame.

3. Question

Answer

Let the speed of rod is v , the length is given by

CLASS24

$$l = l_0 \sqrt{1 - \left(\frac{v^2}{c^2}\right)}$$

Where, l_0 = Length of rod at rest

c = Speed of light $\gg v$

Now assume that the observer and the rod are moving with the speed v in same direction. The length of the rod is given by,

$$l = l_0 \sqrt{1 - \left(\frac{(v - [-v])^2}{c^2}\right)}$$

$$l = l_0 \sqrt{1 - \left(\frac{4v^2}{c^2}\right)}$$

Thus, it is true that an object may be regarded to be at rest or in motion depending on the frame of reference chosen to view the object.

4. Question

Answer

The relativistic mass of the particle is,

$$m = \frac{m_0}{\sqrt{1 - \left(\frac{v^2}{c^2}\right)}}$$

The gravitational force of Earth is given by,

$$F = \frac{GMm_0}{r^2}$$

$$F = \frac{GMm_0}{r^2 \sqrt{1 - \left(\frac{v^2}{c^2}\right)}}$$

Thus, both mass and the gravitational attraction depends on the particle's speed.

5. Question

Answer

CLASS24

There are two ways to measure the distance between the Earth and t which are using meter scale and using light pulse.

In the first method the distance is measured using meter scale, the length gets contracted in moving frame. Thus, the length measured is smaller.

In second method, the time difference between emission of light pulse and reception is noted. So, the distance measured in this case is larger.

Thus, the distance measured is either smaller or larger than the actual distance.

Objective I

1. Question

Answer

The magnitude of linear momentum is given by,

$$P=mv$$

The relativistic mass of the particle is,

$$m = \frac{m_o}{\sqrt{1 - \left(\frac{v^2}{c^2}\right)}}$$

Thus, magnitude of linear momentum is,

$$p = \frac{m_o v}{\sqrt{1 - \left(\frac{v^2}{c^2}\right)}}$$

This value of the linear momentum does no match with any option.

Thus, correct option is D

2. Question

Answer

CLASS24

As the mass is at rest, its value is not changed when observed with the observer moving with the velocity v .

Thus, correct option is C.

3. Question

Answer

The new length is given by,

$$l = l_0 \sqrt{1 - \left(\frac{v^2}{c^2}\right)}$$

Where, l_0 = Length of rod at rest

c = Speed of light $\gg v$

As $v \ll c$

$$\therefore \sqrt{1 - \left(\frac{v^2}{c^2}\right)} < 1$$

Thus, $l < l_0$

Consider the rod at rest and observer moves with the speed of v then the new length is given by,

CLASS24

$$l = l_0 \sqrt{1 - \left(\frac{(-v)^2}{c^2}\right)}$$

$$l = l_0 \sqrt{1 - \left(\frac{(v)^2}{c^2}\right)}$$

Where, l_0 = Length of rod at rest

c = Speed of light $\gg v$

As $v \ll c$

$$\therefore \sqrt{1 - \left(\frac{v^2}{c^2}\right)} < 1$$

Thus, $l < l_0$

Thus, the length will be reduced in both the cases.

So, correct option is C.

4. Question

Answer

The new length is given by,

$$l = l_0 \sqrt{1 - \left(\frac{v^2}{c^2}\right)}$$

Where, l_0 = Length of rod at rest

c = Speed of light $\gg v$

Consider the rod at rest and observer moves with the speed of v parallel to the measured length then the new length is given by,

CLASS24

$$l = l_0 \sqrt{1 - \left(\frac{(-v)^2}{c^2}\right)}$$

$$l = l_0 \sqrt{1 - \left(\frac{(v)^2}{c^2}\right)}$$

Where, l_0 = Length of rod at rest

c = Speed of light $\gg v$

Consider the rod at rest and observer moves with the speed of v opposite to the measured length then the new length is given by,

$$l = l_0 \sqrt{1 - \left(\frac{(v - [-v])^2}{c^2}\right)}$$

$$l = l_0 \sqrt{1 - \left(\frac{4v^2}{c^2}\right)}$$

As $v \ll c$

$$\therefore \sqrt{1 - \left(\frac{v^2}{c^2}\right)} < 1$$

Thus, $l < l_0$

Thus, the length will be minimum for the rod moving in the opposite direction.

So, correct option is B.

5. Question

Answer

The magnitude of linear momentum is given by,

$$P=mv$$

The relativistic mass of the particle is,

$$m = \frac{m_0}{\sqrt{1 - \left(\frac{v^2}{c^2}\right)}}$$

Thus, magnitude of linear momentum is,

$$P = \frac{m_0 v}{\sqrt{1 - \left(\frac{v^2}{c^2}\right)}}$$

$$P = m_0 v \left(1 - \frac{v^2}{c^2}\right)^{-\frac{1}{2}}$$

Using binomial expansion, we get,

$$P = m_0 v + \frac{m_0 v^3}{2c^2}$$

When the speed of the particle is doubled then,

$$p' = \frac{m_0 2v}{\sqrt{1 - \left(\frac{4v^2}{c^2}\right)}}$$

$$P = 2m_0 v + \frac{4m_0 v^3}{c^2}$$

Thus, the new value of linear momentum is more than double.

So, correct option is B.

6. Question

Answer

When the force acting on the particle is constant, the speed of the particle increases due to the force.

The relativistic mass of the particle is,

$$m = \frac{m_0}{\sqrt{1 - \left(\frac{v^2}{c^2}\right)}}$$

So, the mass of the particle increases, this results in decreasing the acceleration gradually.

Thus, correct option is B.

7. Question

Answer

The charged particle is projected at a very high speed perpendicular to the uniform magnetic field. The mass will increase using the relativistic relation.

The relativistic mass of the particle is,

$$m = \frac{m_0}{\sqrt{1 - \left(\frac{v^2}{c^2}\right)}}$$

So, its radius will increase. Thus, correct option is B.

Objective II

1. Question

Answer

Statement (a) is correct because the equation of special relativity is only applicable when the speed of object is comparable to the speed of light. Newton's

equations of motion are used for small speeds.

CLASS24

When the body is travelling with the relativistic speed, then

mass becomes $m' = \gamma m$, length becomes $L' = \frac{L}{\gamma}$, change in time becomes

$$\Delta T' = \gamma \Delta T \text{ where } \gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$\Rightarrow \gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} = \left(1 - \frac{v^2}{c^2}\right)^{-\frac{1}{2}} = 1 + \frac{v^2}{2c^2} + \dots = \text{sum} \ll 1 \text{ when } v < c.$$

Still the sum of the series will be greater than 0

Hence, non-relativistic equations in which γ factor is taken to be exactly 1 never give exact results.

Option (B) and (D) are correct

2. Question

Answer

If the speed of rod is relativistic speed then, its mass will be

$$m' = \gamma m = \frac{m}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$\Rightarrow \gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} = \left(1 - \frac{v^2}{c^2}\right)^{-\frac{1}{2}} = 1 + \frac{v^2}{2c^2} + \dots > 1 \text{ as } v < c$$

And its length will be,

$$L' = \frac{L}{\gamma} = L \sqrt{1 - \frac{v^2}{c^2}}$$

When speed v is doubled,

$$\gamma' = \frac{1}{\sqrt{1 - \frac{4v^2}{c^2}}}$$

$$\Rightarrow \gamma' = \left(1 - \frac{4v^2}{c^2}\right)^{-\frac{1}{2}} = 1 + \frac{2v^2}{c^2} + \dots > 2\gamma$$

and $m' = \gamma' m > 2\gamma m$, $L' = \frac{L}{\gamma'} < \frac{L}{2\gamma}$

Hence, mass will be increased and it will be more than double. Length will decrease but not exactly half of the original length.

Option (C) and (D) are correct.

3. Question


Answer

The two events occur at the same time at points A and B in rest frame. If they occur simultaneously in a frame moving with respect to the lab, the frame must be moving in the direction perpendicular to AB.

Option (b) is correct.

4. Question

Answer



If the electron is moving with very high-speed v , then its mass will increase by the factor γ i.e. $m' = \gamma m$, where $\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$,

CLASS24

Momentum, $p = m'v = \gamma mv$

Kinetic energy $k = \frac{1}{2} m'v^2 = \frac{1}{2} \gamma mv^2$

When the speed of electron is maximum i.e. $v=c$, then $\gamma = \infty$, also $m', p, k = \infty$

Therefore, the upper limit of speed is always less than c but mass, momentum and kinetic energy do not have any upper limit.

Option (c) is correct.

5. Question

Answer

the rest length of the rod = L

Relativistic speed = v

Length contraction = L'

$$L' = \frac{L}{\gamma} = L \sqrt{1 - \frac{v^2}{c^2}}$$

$$\Rightarrow \gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

If the rod is moving with relativistic speed, then γ will be approximately equal to 1. This means that the length of the rod L' will also be almost equal to L . However, the length of rod in rest frame L , can be more than L' depending upon the frame of observer.

Option (B) and (c) are correct

6. Question

Answer

if the rod is moving with relativistic speed v , then mass will be given by,

$$m' = \gamma m = \frac{m}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$\text{here, } \gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

thus, the mass will increase by the factor of γ

option (A) is correct

Exercises

1. Question

Answer

The house of yogi is 1000km away from his guru in Himalayan cave. The velocity of travelling of the thoughts should be maximum so that it can reach to his guru in minimum time. The maximum velocity that can be attained is speed of light i.e. 3×10^8 m/sec.

$$\text{Distance} = 1000 \text{ km} = 10^6 \text{ m}$$

$$\text{Velocity} = 3 \times 10^8 \text{ m/sec}$$

$$\text{Time} = \frac{\text{Distance}}{\text{velocity}} = \frac{10^6}{3 \times 10^8} = \frac{1}{300} \text{ sec}$$

2. Question

- (a) the traveler and
- (b) a ground observer

Answer

(a) The traveler is traveling in the same frame in which suitcase is travelling. So, the dimensions of the suitcase for the traveler will remain same.

(b) The observer who is on the ground will not see the same dimensions of the suitcase. It is because the frame of reference of observer and the suitcase is different. The suitcase is in the frame of reference which is moving with the speed of $0.6c$. Since, the suitcase is placed with its length along the train's velocity, the changes will be observed in the length only.

Length of the suitcase= 50cm

Breadth = 25cm

Height = 10cm

Velocity= $0.6c$

$$L' = L \sqrt{1 - \frac{v^2}{c^2}} = 50 \sqrt{1 - \frac{(0.6c)^2}{c^2}} = 50 \sqrt{1 - 0.36} = 50 \times 0.8 = 40\text{cm}$$

Therefore, the length observed by the observer on the ground will be less than the original length of the suitcase. The dimensions will be 40cm x 25cm x 10cm.

3. Question

Answer

Whenever the object moves with the speed comparable to the speed of light, the length of the object contracts. So, let us consider the following cases when the speed of rod is comparable to the speed of light.

(a) length of the rod= 1m

Speed of the rod= $3 \times 10^5 \text{ m s}^{-1}$

Using the relativistic relation of length contraction

$$L' = L \sqrt{1 - \frac{v^2}{c^2}} = 1 \sqrt{1 - \frac{9 \times 10^{10}}{9 \times 10^{16}}} = \sqrt{1 - 10^{-6}} = 0.99999995 \quad \text{CLASS24}$$

The length observed will be 0.99999995m

(b) length = 1m

$$\text{Speed} = 3 \times 10^6 \text{ m s}^{-1}$$

$$L' = L \sqrt{1 - \frac{v^2}{c^2}} = 1 \sqrt{1 - \frac{9 \times 10^{12}}{9 \times 10^{16}}} = \sqrt{1 - 10^{-4}} = 0.99995\text{m}$$

(c) length = 1m

$$\text{Speed} = 3 \times 10^7 \text{ m s}^{-1}$$

$$L' = L \sqrt{1 - \frac{v^2}{c^2}} = 1 \sqrt{1 - \frac{9 \times 10^{14}}{9 \times 10^{16}}} = \sqrt{1 - 10^{-2}} = 0.9949\text{m}$$

4. Question

Answer

The velocity of the train, $v = 0.6c$

Time taken by the train to pass the man standing on platform, $t = 1\text{sec}$

(a) the length of the train as seen by the person, $L' = vt = 0.6 \times 3 \times 10^8 = 1.8 \times 10^8\text{m}$

(b) the rest length of the train L

$$L' = L \sqrt{1 - \frac{v^2}{c^2}}$$

$$1.8 \times 10^8 = L \sqrt{1 - \frac{(0.6c)^2}{c^2}}$$

$$L = \frac{1.8 \times 10^8}{0.8} = 2.25 \times 10^8 \text{ m/sec}$$

5. Question

Answer

CLASS24

The field becomes square in the plane frame only when its length and breadth become equal. Originally, the field is in rectangular shape, with length = 100m and breadth = 50 m. In order to make the rectangular field into square field, either the breadth can be increased or length can be reduced. But we know that with the speed comparable to the speed of light, the object's length contracts. Hence, the only possible way to make the rectangular field into square field is that the plane must travel along the length of the field with the speed comparable to the speed of light.

$$L' = L \sqrt{1 - \frac{v^2}{c^2}}$$

$$\Rightarrow 50 = 100 \sqrt{1 - \frac{v^2}{c^2}}$$

$$\Rightarrow \frac{1}{2} = \sqrt{1 - \frac{v^2}{c^2}}$$

$$\Rightarrow v = 0.866c$$

6. Question

Answer

The distance between Patna and Delhi = 1000km

Speed of the train = $360 \text{ km h}^{-1} = 100 \text{ m/s}$

(a) Distance between the two cities in the train frame d

$$D' = D \sqrt{1 - \frac{v^2}{c^2}}$$

$$\Rightarrow D' = 10^6 \sqrt{1 - \frac{10^4}{9 \times 10^{16}}}$$

$$\Rightarrow D' = 10^9$$

(b) time elapses in the train frame

$$\Delta t = \frac{\Delta L}{v}$$

To solve for ΔL

$$L' = L \sqrt{1 - \frac{v^2}{c^2}}$$

$$L' = 10^6 \sqrt{1 - \frac{100^2}{c^2}}$$

$$\Rightarrow L' = 56 \times 10^{-9} \text{ m}$$

Change in length $\Delta L = 56 \text{ nm}$

therefore, the distance between Patna and Delhi in the train frame is 56 nm less than 1000 km.

now, actual time taken by train = time = distance / speed

$$t = \frac{1000 \times 10^3}{100} = 10^4 \text{ sec}$$

$$\text{change in time, } \Delta T = \frac{\Delta L}{v} = \frac{56 \times 10^{-9}}{100} = 0.56 \text{ ns}$$

So, the time lapse in the train between Patna and Delhi will be 0.56 ns less than 10^4 sec

7. Question

Answer

Speed of the car = $180 \text{ km h}^{-1} = 50 \text{ m/sec}$

Time taken by car to reach from station A to station B = 10 hours

The distance travelled by person in a car $L' = \text{speed} \times \text{time} = 180 \times 10 = 1800 \text{ km}$

$$L' = L \sqrt{1 - \frac{v^2}{c^2}}$$

$$1800 = L \sqrt{1 - \frac{(180)^2}{9 \times 10^{16}}}$$

$$\Rightarrow L = 1800 + 25 \times 10^{-9}$$

So, the rest distance between the two station is 25nm more than 1800km.

(b) time taken by car to cover the distance in the road= distance/ speed

$$\Rightarrow \frac{1.8 \times 10^6 + 25 \times 10^{-9}}{50}$$

$$\Rightarrow 0.36 \times 10^5 + 5 \times 10^{-8} = 10 \text{ hours } 0.5 \text{ ns}$$

8. Question

Answer

The speed of spaceship = $5c/13$

(a) consecutive birthday celebration means the time period $t = 1$ year.

But the time interval observed by the person in spacecraft will be

$$t' = \frac{t}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$\Rightarrow t' = \frac{1}{\sqrt{1 - \frac{(5c/13)^2}{c^2}}}$$

$$\Rightarrow t' = \frac{1}{\sqrt{1 - \frac{25}{169}}}$$

$$\Rightarrow t' = \frac{13}{12} \text{ years}$$

Thus, the time interval observed by the person will be more than one year i.e. 1.08 years (approx.)

CLASS24

(b) The person on Earth will also calculate the same time interval because for him also, the person in spacecraft is travelling with same speed.

9. Question

Answer

Both the stations are in ground frame; hence the station clocks will record the proper time interval. But the clocks in train will record improper time because they are at different places travelling with different speed. The proper time interval ΔT is less than improper i.e. $\Delta T' = \gamma \Delta T$. Hence, in case (a) In the train going from Howrah to Delhi, the baby born in Delhi will be elder. (b) In the train going from Delhi to Howrah, Howrah baby will be elder.

10. Question

Answer

Since, the frame is moving, the clocks will not record the synchronized time. The clock at the rear end leads the clock which is at the other end by $L v/c^2$ where, L is the rest separation between the clocks and v is the speed of the moving frame i.e. train. Thus, the baby born adjacent to the guard cell will be elder.

11. Question

Answer

velocity of Swarglok with respect to Earth = $0.9999c$

According to the Earth' frame the velocity will be

$$v' = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$\Rightarrow v' = \frac{1}{\sqrt{1 - \frac{(0.9999c)^2}{c^2}}} = \frac{1}{0.01414} = 70.712$$

Let one day on Earth = ΔT

One day on heaven = $\Delta T'$

$$\Delta T' = v \Delta T$$

$$\Rightarrow \Delta T' = 70.71 \text{ days in heaven}$$

12. Question

Answer

As the Lorentz transformation suggests there will be time dilation for a moving object with respect to rest frame which is given by γT_0

Where $\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$, v is velocity of moving object and c is speed of light in vacuum i.e. $3 \times 10^8 \text{ m/s}$

And T_0 is time in rest frame.

ATQ, $v = 60\% \text{ of } c$ i.e. $\frac{v}{c} = \frac{3}{5}$

$$\therefore \gamma = \frac{1}{\sqrt{1 - \left(\frac{3}{5}\right)^2}} = \frac{5}{4}$$

\therefore The person will live $\frac{5}{4} \times 100 \text{ years} = 125 \text{ years}$ in the spaceship with respect to earth frame.

13. Question

Answer

As the Lorentz transformation suggests there will be time dilation for a moving object with respect to rest frame which is given by γT_0

CLASS24

Where $\gamma = \frac{1}{\sqrt{1-v^2/c^2}}$, v is velocity of moving object and c is speed of light in vacuum i.e. $3 \times 10^8 \text{ m/s}$

And T_0 is time in rest frame.

$$\text{ATQ, } \frac{v}{c} = \frac{4}{5}$$

$$\therefore \gamma = \frac{1}{\sqrt{1 - \left(\frac{4}{5}\right)^2}} = \frac{5}{3}$$

\therefore Time taken by the bulb to switch off and switch on with respect to spaceship will be $\frac{5}{3} \times 1\text{s}$

\therefore Frequency will be $\frac{1}{T} = \frac{3}{5} \text{ Hz}$

14. Question

Answer

Time taken by the car to reach the tower B from tower A $= \frac{1000}{100} = 10 \text{ Hour}$.

As the frame of wrist watch is moving \therefore the wrist watch will lag by $\gamma T_0 - T_0$.

Where $\gamma = \frac{1}{\sqrt{1-v^2/c^2}}$, v is velocity of moving object and c is speed of light in vacuum i.e. $3 \times 10^8 \text{ m/s}$

And T_0 is time in rest frame .

$$V = 100 \text{ Km/hr} = 100 \times \frac{5}{18} \text{ m/s} = 27.78 \text{ m/s}$$

$\frac{v}{c} = \frac{27.78}{3 \times 10^8}$ as $\frac{v}{c} \ll 1$ using binomial expansion we know we can write

$$(1 + x)^n = 1 + nx \text{ when } x \ll 1$$

$$\text{Similarly, } \gamma = \frac{1}{\sqrt{1 - \left(\frac{27.78}{3 \times 10^8}\right)^2}} = \left(1 - \left(\frac{27.78}{3 \times 10^8}\right)^2\right)^{-\frac{1}{2}} = 1 + \frac{1}{2} \times 86.0254 \times 10^{-16}$$

$$\therefore T_o(\gamma - 1) = 10 \text{ hour} \times 43 \times 10^{-16} = 1.548 \times 10^{-12} \text{ s}$$

Since 1 hour = 3600 s

CLASS24

And as we can see the calculated value is too small to make drastic affect that's why we don't face any trouble in real life situation.

15. Question

Answer

As we know the moving object appears to be contracted with the earth aur rest frame that's because of Lorentz contraction known as length contraction which is

given by $l = \frac{l_o}{\gamma}$

Where $\gamma = \frac{1}{\sqrt{1 - v^2/c^2}}$, V is velocity of moving object and C is speed of light in vacuum i.e. $3 \times 10^8 \text{ m/s}$

And l_o is length at rest .

As volume always include power of 3 weather it is cube sphere we will take a simple case of cube where all of its length is given by l_o at rest

\therefore Volume at rest is $= l_o^3$

ATQ Volume seems to be half at certain speed

$$\therefore l^3 = \frac{l_o^3}{2}$$

$$\Rightarrow l = \frac{l_o}{\sqrt[3]{2}} = 0.794l_o$$

$$\therefore \frac{l_o}{\gamma} = 0.794l_o$$

$$\Rightarrow \gamma = 1.26$$

And $1.26 = \frac{1}{\sqrt{1 - v^2/c^2}}$

$$\Rightarrow \sqrt{1 - v^2/c^2} = 0.63$$

$$\Rightarrow 0.604 \times c^2 = v^2$$

$$\therefore v = 0.78c$$

16. Question

Answer

CLASS24

a) We know time taken = $\frac{\text{Distance traversed}}{\text{Speed of the particle}}$

$$\therefore \text{life of particle will be} = \frac{1 \times 10^{-2} \text{ m}}{0.995 \times 3 \times 10^8 \text{ m/s}} = 3.35 \times 10^{-11} \text{ s}$$

b) In the frame of particle there will be length contraction or we can say time dilation we can solve the question with anyone of the approach.

As the Lorentz transformation suggests there will be time dilation for a moving object with respect to rest frame which is given by γT_0

Where $\gamma = \frac{1}{\sqrt{1 - v^2/c^2}}$, v is velocity of moving object and c is speed of light in vacuum i.e. $3 \times 10^8 \text{ m/s}$

And T_0 is time in rest frame.

$\therefore T_0$ was found in previous case as $3.35 \times 10^{-11} \text{ s}$

$$\therefore \text{Life of particle in particle frame will be} = T_0 \times \frac{1}{\sqrt{1 - v^2/c^2}}$$

$$\Rightarrow 3.35 \times 10^{-11} \text{ s} \times \frac{1}{\sqrt{1 - (0.995c)^2/c^2}} = 3.35 \times 10^{-11} \text{ s} \times 10$$

$$= 3.35 \times 10^{-10} \text{ s}$$

17. Question

Answer

There is a physical significance of special theory of relativity that Energy and mass are interchangeable and that is given by Sir Einstein famous equation $E=mc^2$

∴ energy possessed by the compressed spring as potential energy is due to loss of the mass

CLASS24

$$\therefore \frac{1}{2} kx^2 = mC^2$$

Given:

$$k = 500 \text{ N/m}$$

$$x = 1 \text{ cm} = 10^{-2} \text{ m}$$

$$\Rightarrow \frac{1}{2} \times 500 \times (10^{-2})^2 = m \times 9 \times 10^{16}$$

$$\therefore m = 2.78 \times 10^{-19} \text{ Kg}$$

So, the change in mass of spring will be $2.78 \times 10^{-19} \text{ Kg}$

18. Question

Answer

There is a physical significance of special theory of relativity that Energy and mass are interchangeable and that is given by Sir Einstein famous equation $E=mc^2$

∴ Heat energy given will be converted in the mass

$$\Delta Q = mc\Delta T = \Delta mC^2$$

Given:

$$c = \text{Specific heat capacity of water} = 4200 \text{ J kg}^{-1} \text{ K}^{-1}$$

$$m = 1 \text{ kg}$$

$$\Delta T = 373 \text{ K} - 273 \text{ K} = 100 \text{ K}$$

$$\therefore 1 \times 4200 \times 100 = \Delta m \times 9 \times 10^{16}$$

$$\Rightarrow \Delta m = \frac{4.2 \times 10^5}{9 \times 10^{16}} = 4.7 \times 10^{-12} \text{ Kg}$$

19. Question

Answer

There is a physical significance of special theory of relativity that Energy and mass are interchangeable and that is given by Sir Einstein famous equation

CLASS24

∴ Heat energy extracted will result in loss of the mass

$$\Delta Q = C_v \Delta T = \Delta m C^2$$

Given:

$$C_v = \frac{3}{2}R \text{ where } R = 8.314 \text{ J/K.mol}$$

$$\Delta T = 10K$$

$$\therefore \frac{3}{2} \times 8.314 \times 10 = \Delta m \times 9 \times 10^{16}$$

$$\Rightarrow \Delta m = 13.86 \times 10^{-16} Kg$$

20. Question

Answer

There is a physical significance of special theory of relativity that Energy and mass are interchangeable and that is given by Sir Einstein famous equation $E=mc^2$

∴ Gain in kinetic energy by the boy is possessed by the mass gain for the boy.

$$\therefore \frac{1}{2} \times m \times V^2 = \Delta m C^2$$

Given:

$$V=12 \text{ km/h} = 12 \times \frac{5}{18} \text{ m/s}$$

$$\therefore \frac{1}{2} \times m \times \left(12 \times \frac{5}{18}\right)^2 = \Delta m \times 9 \times 10^{16}$$

$$\Rightarrow \frac{\Delta m}{m} = \frac{1}{2} \times \frac{1}{9 \times 10^{16}} \times \left(12 \times \frac{5}{18}\right)^2$$

$$\Rightarrow \frac{\Delta m}{m} = 6.17 \times 10^{-17}$$

21. Question

Answer

There is a physical significance of special theory of relativity that Energy and mass are interchangeable and that is given by Sir Einstein famous equation

CLASS24

∴ Loss in the mass due to continuous illumination of bulb

ATQ, the bulb illuminates for 1 year i.e $365 \times 24 \times 3600$ s with power of 1 W or 1 J/s.

Total energy dissipated by the bulb in the form of light energy will

$$\text{be } 100 \times 365 \times 24 \times 3600 \text{ J} = 3.15 \times 10^9 \text{ J}$$

$$\therefore 3.15 \times 10^9 \text{ J} = \Delta m \times 9 \times 10^{16}$$

$$\Rightarrow \Delta m = 3.5 \times 10^{-8} \text{ kg}$$

22. Question

Answer

a)

The amount of energy radiated by the sun in $1 \text{ m}^2 = 1400 \text{ W}$

As the energy radiated is dependent on the surface area it covers which is given by $4 \times \pi \times r^2$

∴ Energy loosed by the sun till it reaches earth in 1 second is given by

$$4 \times \pi \times (1.5 \times 10^{11} \text{ m})^2 \times 1400 = 3.96 \times 10^{26} \text{ J}$$

So, the mass loosed by the sun in 1 second is $\frac{3.96 \times 10^{26}}{9 \times 10^{16}} = 4.4 \times 10^9 \text{ Kg}$

b)

As the rate of decay of the mass is constant it will last up to

$$\frac{\text{Total mass of sun}}{\text{Rate at which mass is decaying}} = \frac{2 \times 10^{30} \text{ Kg}}{4.4 \times \frac{10^9 \text{ Kg}}{\text{s}}} = 4.55 \times 10^{10} \text{ s}$$

$$\text{In years } \frac{4.55 \times 10^{10} \text{ s}}{365 \times 24 \times 3600 \text{ s}} = 1.44 \times 10^3 \text{ Years}$$

23. Question

Answer

As we know electron and proton has same mass i.e. $9.1 \times 10^{-31} \text{Kg}$

CLASS24

As they annihilate and produces gamma particle with some energy

$$\Delta m = 2 \times 9.1 \times 10^{-31} \text{Kg}$$

There is a physical significance of special theory of relativity that Energy and mass are interchangeable and that is given by Sir Einstein famous equation $E = \Delta m C^2$

$$\begin{aligned} \therefore \text{Energy of gamma particle} &= 2 \times 9.1 \times 10^{-31} \text{Kg} \times 9 \times 10^{16} \frac{\text{m}}{\text{s}} \\ &= 1.66 \times 10^{-13} \text{J} \end{aligned}$$

24. Question

Answer

As we know the moving object appears to be heavier in the moving frame that's because of Lorentz transformation known as apparent mass which is given by

$$m = m_o \gamma$$

Where $\gamma = \frac{1}{\sqrt{1 - v^2/c^2}}$, v is velocity of moving object and C is speed of light in vacuum i.e. $3 \times 10^8 \text{ m/s}$

And m_o is mass at rest

$$\therefore \gamma = \frac{1}{\sqrt{1 - (0.8C)^2/C^2}} = \frac{1}{0.6}$$

$$\therefore \text{Apparent mass} = \frac{9.1 \times 10^{-31} \text{Kg}}{0.6} = 1.52 \times 10^{-30} \text{Kg}$$

Kinetic energy gained by the electron will be due to change in mass of electron.

$$(m - m_o)C^2 = (15.2 - 9.1) \times 10^{-31} \times 9 \times 10^{16} = 5.45 \times 10^{-15} \text{J}$$

Momentum of electron will be is equal to = Velocity \times apparent mass

$$\Rightarrow 1.52 \times 10^{-30} \text{Kg} \times 0.8 \times 3 \times 10^8 \text{ m/s} = 3.65 \times 10^{-22} \text{ kg.m/s}$$

25. Question

Answer

a) As we know the moving object appears to be heavier in the moving frame because of Lorentz transformation known as apparent mass which is given by

$$m = m_0 \gamma$$

Where $\gamma = \frac{1}{\sqrt{1 - v^2/c^2}}$, v is velocity of moving object and c is speed of light in vacuum i.e. $3 \times 10^8 \text{ m/s}$

And m_0 is mass at rest

$$\therefore \gamma = \frac{1}{\sqrt{1 - (0.6c)^2/c^2}} = \frac{1}{0.8}$$

$$\therefore \text{Apparent mass} = \frac{9.1 \times 10^{-31} \text{ Kg}}{0.8} = 11.375 \times 10^{-31} \text{ Kg}$$

Kinetic energy gained by the electron will be due to change in mass of electron.

$$(m - m_0)c^2 = (11.375 - 9.1) \times 10^{-31} \times 9 \times 10^{16} = 20.475 \times 10^{-15} \text{ J}$$

As the gain in kinetic energy is given by charge \times potential difference applied i.e. $q \times V$

$$\text{As } q = 1.6 \times 10^{-19} \text{ C}$$

$$\therefore 1.6 \times 10^{-19} \text{ C} \times V = 20.475 \times 10^{-15} \text{ J}$$

$$\Rightarrow V = 12.8 \times 10^4 \text{ V}$$

b) Similarly for $0.9c$

$$\gamma = \frac{1}{\sqrt{1 - (0.9c)^2/c^2}} = \frac{1}{0.44}$$

$$\therefore \text{Apparent mass} = \frac{9.1 \times 10^{-31} \text{ Kg}}{0.44} = 20.68 \times 10^{-31} \text{ Kg}$$

Kinetic energy gained by the electron will be due to change in mass of electron.

$$(m - m_0)c^2 = (20.68 - 9.1) \times 10^{-31} \times 9 \times 10^{16} = 1 \times 10^{-13} \text{ J}$$

As the gain in kinetic energy is given by charge \times potential difference applied i.e. $q \times V$

$$\text{As } q = 1.6 \times 10^{-19} \text{ C}$$

$$\therefore 1.6 \times 10^{-19} \text{ C} \times V = 1 \times 10^{-13} \text{ J}$$

$$\Rightarrow V = 6.25 \times 10^5 V$$

c) Similarly for 0.99c

$$\gamma = \frac{1}{\sqrt{1 - (0.99c)^2/c^2}} = \frac{1}{0.1}$$

$$\therefore \text{Apparent mass} = \frac{9.1 \times 10^{-31} Kg}{0.1} = 9.1 \times 10^{-30} Kg$$

Kinetic energy gained by the electron will be due to change in mass of electron.

$$(m - m_0)C^2 = (91 - 9.1) \times 10^{-31} \times 9 \times 10^{16} = 7.37 \times 10^{-13} J$$

As the gain in kinetic energy is given by charge \times potential difference applied i.e. $q \times V$

$$\text{As } q = 1.6 \times 10^{-19} C$$

$$\therefore 1.6 \times 10^{-19} C \times V = 7.37 \times 10^{-13} J$$

$$\Rightarrow V = 4.6 \times 10^6 V$$

26. Question

Answer

We can't deal the question with classical mechanics approach as $\frac{1}{2} \times m_e \times v^2$ as when you calculate v with this approach it will come out to be more than speed of light in vacuum i.e. $3 \times 10^8 \text{ m/s}$ that violates the special theory of relativity.

So we have to solve these types of question taking relativistic approach.

As we know the gain in Kinetic energy is due to change in mass

As we know the moving object appears to be heavier in the moving frame that's because of Lorentz transformation known as apparent mass which is given by

$$m = m_0 \gamma$$

Where $\gamma = \frac{1}{\sqrt{1 - v^2/c^2}}$, v is velocity of moving object and C is speed of light in vacuum i.e. $3 \times 10^8 \text{ m/s}$

And m_0 is mass at rest i.e. $= 9.1 \times 10^{-31} Kg$

$$K.E = m_0(\gamma - 1)C^2$$

$$\therefore m_0 \times C^2 = 9.1 \times 10^{-31} \times 9 \times 10^{16} = 8.19 \times 10^{-14} \text{kg m}^2/\text{s}^2$$

a) So 10 for 1 eV i.e. $1.6 \times 10^{-19} \text{J}$

$$8.19 \times 10^{-14} \text{kg m}^2/\text{s}^2 \times (\gamma - 1) = 1.6 \times 10^{-19} \text{J}$$

$$\Rightarrow \gamma - 1 = 1.95 \times 10^{-6}$$

$$\Rightarrow \gamma = 1 + 1.95 \times 10^{-6}$$

$$\text{As } \gamma = \frac{1}{\sqrt{1 - V^2/C^2}}$$

Squaring both side and using binomial as we know using binomial expansion we know we can write

$$(1 + x)^n = 1 + nx \text{ when } x \ll 1$$

$$\therefore 1 + 3.9 \times 10^{-6} = \frac{1}{1 - V^2/C^2}$$

$$\Rightarrow 1 - V^2/C^2 = \frac{1}{1 + 3.9 \times 10^{-6}}$$

$$\Rightarrow V^2 = C^2 \times 3.9 \times 10^{-6}$$

$$V = 6 \times 10^5 \text{ m/s}$$

b) Similarly for 10KeV i.e. $1.6 \times 10^{-15} \text{J}$

$$8.19 \times 10^{-14} \text{kg m}^2/\text{s}^2 \times (\gamma - 1) = 1.6 \times 10^{-15} \text{J}$$

$$\Rightarrow \gamma - 1 = 1.95 \times 10^{-2}$$

$$\Rightarrow \gamma = 1 + 1.95 \times 10^{-2}$$

$$\text{As } \gamma = \frac{1}{\sqrt{1 - V^2/C^2}}$$

Squaring both side and using binomial as we know using binomial expansion we know we can write

$$(1 + x)^n = 1 + nx \text{ when } x \ll 1$$

$$\therefore 1 + = \frac{1}{1 - V^2/C^2}$$

$$\Rightarrow 1 - V^2/C^2 = \frac{1}{1 + 3.8 \times 10^{-2}}$$

$$\Rightarrow V^2 = C^2 \times 3.9 \times 10^{-2}$$

$$V = 6 \times 10^7 \text{ m/s}$$

c) Similarly for 10KeV i.e. $1.6 \times 10^{-12} \text{ J}$

$$8.19 \times 10^{-14} \text{ kg } \frac{\text{m}^2}{\text{s}^2} \times (\gamma - 1) = 1.6 \times 10^{-12} \text{ J}$$

$$\Rightarrow \gamma - 1 = 19.5$$

$$\Rightarrow \gamma = 20.5$$

$$\text{As } \gamma = \frac{1}{\sqrt{1 - V^2/C^2}}$$

$$\therefore 20.5 = \frac{1}{1 - V^2/C^2}$$

$$\Rightarrow 1 - V^2/C^2 = 0.0024$$

$$\Rightarrow V^2 = C^2 \times 0.9976$$

$$\therefore V = 0.998C$$

27. Question

Answer

As we know that Kinetic energy gained by the electron is due to change in mass of electron.

ATQ mass becomes double $\therefore \Delta m = 2m_0 - m_0$

And m_0 is mass at rest i.e. $= 9.1 \times 10^{-31} \text{ Kg}$

$$\text{As } E = \text{K.E} = \Delta m C^2 = m_0 \times 9 \times 10^{16} = 9.1 \times 10^{-31} \times 9 \times 10^{16} = 8.2 \times 10^{-14} \text{ J}$$

28. Question

Answer

As we know that Kinetic energy gained by the electron is due to change in mass of electron.

$$\text{As } E = \text{K.E} = \Delta m C^2$$

As we know relativistic Energy is always more than non-relativistic kinetic energy because Rest energy is always less than apparent mass energy

CLASS24

∴ The relativistic value of Kinetic energy will be $\frac{101}{200} \times m_0 \times V^2$

As we know the moving object appears to be heavier in the moving frame that's because of Lorentz transformation known as apparent mass which is given by

$$m = m_0 \gamma$$

Where $\gamma = \frac{1}{\sqrt{1 - V^2/C^2}}$, V is velocity of moving object and C is speed of light in vacuum i.e. $3 \times 10^8 \text{ m/s}$

$$\text{K.E} = m_0(\gamma - 1)C^2 = \frac{101}{200} \times m_0 \times V^2$$

$$\Rightarrow \frac{1}{\sqrt{1 - V^2/C^2}} - 1 = \frac{101}{200} \times m_0 \times \frac{V^2}{C^2}$$

$$\text{Let } \frac{V^2}{C^2} = K$$

$$\therefore \frac{1}{\sqrt{1 - K}} - 1 = \frac{101}{200} \times K$$

$$\Rightarrow \frac{1}{\sqrt{1 - K}} = \frac{301}{200} \times K$$

Squaring both side we get

$$\Rightarrow \frac{1}{1 - K} = 2.265 \times K^2$$

$$\Rightarrow 1 = 2.265K^2 - 2.265K^3$$

As $K = \frac{V^2}{C^2}$, $K^3 = \frac{V^6}{C^6}$ which is $\ll 1$ ∴ we can neglect K^3 term.

$$\Rightarrow K^2 = \frac{1}{2.265} \Rightarrow K = 0.441$$

$$\Rightarrow \frac{V^2}{C^2} = 0.441$$

$$\therefore V = 0.66 C$$